

An Automated Search for Metal-Poor Halo Stars in the Hamburg/ESO Objective-Prism Survey

Norbert Christlieb¹, Lutz Wisotzki & Dieter Reimers

Hamburger Sternwarte, Germany

Thomas Gehren & Johannes Reetz

Universitäts-Sternwarte München, Germany

Timothy C. Beers

Department of Physics & Astronomy, Michigan State University, U.S.A.

Abstract. An automated search for metal-poor stars is carried out in the course of the Hamburg/ESO objective-prism survey (HES), which covers the full southern sky at Galactic latitudes $|b| \gtrsim 30^\circ$. As the HES reaches ~ 1 magnitude deeper and covers areas of the sky which have not been touched by the HK Survey of Beers and collaborators, the total survey volume of the HES represents an increase by a factor of 4.5 compared to the HK Survey. Because of the limited availability of telescope time for spectroscopic follow-up observations, we will focus on a search for the most metal-poor, unevolved stars. We present a simulation study of the HES selection function $P(B_J, T_{\text{eff}}, [\text{Fe}/\text{H}], \log g)$ for metal-poor stars, and results from a first follow-up campaign at the ESO-NTT.

1. Introduction

Metal-poor stars have atmospheres that consist of the most primitive (luminous) matter in the Universe. They offer the possibility to study the formation and early chemical evolution of our Galaxy, and the first epochs of star formation. Moreover, they are of significance for observational cosmology, e.g., by setting a lower limit to the age of the Universe by their ages, and by measurement of the abundances of light element lithium, thought to have been formed in Big Bang nucleosynthesis.

As a result of the interference-filter/objective-prism survey of Beers and collaborators (Beers et al. 1992; hereafter BPS92), the so-called “HK Survey,” the number of known metal-deficient stars has increased dramatically in the last few years. However, as emphasized by Beers (1999, this volume), many interesting projects to be undertaken in the near future require still larger samples of *extremely* ($[\text{Fe}/\text{H}] < -2.5$) metal-poor stars.

¹E-mail: nchristlieb@hs.uni-hamburg.de

Given this fact, but also facing the limited availability of observing time for spectroscopic follow-up observations, we have decided to focus on a sub-sample of the metal-poor stars being present on our plates, namely the most metal-poor, unevolved stars (main-sequence stars, turnoff-stars and subgiants). The restriction of the candidate set to these stars results in a tractable number of candidates per field. In section 3 we show how the selection can be optimized towards this aim.

With the advent of large multi-object spectrographs, e.g., 6dF (Watson 1998), projects involving follow-up observations of a larger number of metal-poor stars should soon be feasible, such as detailed investigation of the metallicity distribution function of the Galactic Halo (Beers 1999, this volume).

2. The Hamburg/ESO Survey

The HES (Wisotzki et al. 1996) is an objective-prism survey primarily targetted for bright QSOs, covering the full southern extragalactic sky ($\sim 10\,000\text{ deg}^2$). It is based on IIIa-J plates which have been taken with the ESO Schmidt telescope and its 4° prism, yielding a wavelength range of $3200\text{ \AA} < \lambda < 5300\text{ \AA}$ and a seeing-limited spectral resolution of $15\text{ \AA}$ at $H\gamma$.

So far 80 % of the plates have been digitised in Hamburg using a PDS 1010G microdensitometer. Scanning and reduction of all plates is expected to be completed by the end of 1998. Apart from the quasar survey work, this data base of digitised moderate-resolution spectra opens the possibility of a fast and efficient exploitation of the *stellar* content of the survey, which includes, among many other interesting classes of objects, the most metal-poor stars.

Because the limiting magnitude of the HES plates ($B \sim 16.5$) is about one magnitude fainter than that of the HK Survey ($B \sim 15.5$), and a large area in the southern extragalactic sky which is not covered by the HK survey is covered by the HES (cf. Fig. 1), the HES has the potential for another significant increase of the number of known metal-poor stars.

3. Selection of Candidates

As a benchmark for a selection of metal-poor stars by automatic classification, a technique which is under development (Christlieb et al. 1997; 1998a,b), we implemented a search for spectra exhibiting a Ca II K line weaker than normal for its colour. Thanks to the *UBV* photometry of the HK survey, it was possible to perform a calibration between $B - V$ and the half-power point of the density distribution of the HES spectra in the range $3900\text{ \AA} < \lambda < 5300\text{ \AA}$ (named **x_hpp2** in the following). The r.m.s. error of the calibration is 0.1 mag, more than sufficient for this purpose.

“Normal” Ca II K line strengths are defined plate-by-plate by a straight line fit of the Ca II K index as a function of **x_hpp2**. The fit is performed for all point-like, overlap-free sources with $S/N > 15$ on one plate, after outliers (e.g., metal-poor stars, but also many plate artifacts) have been removed by an iterative $\kappa\sigma$ -clipping.




Figure 1. Comparison of HK survey and Hamburg/ESO survey volumes, assuming limiting magnitudes of $B_J = 15.5$ and $B_J = 16.5$, respectively. By the HES the total survey volume for metal-poor stars can be increased by a factor of 4.5 compared to the HK survey alone.

3.1. A Simulation Study of the Metal-Poor Star Selection Function

In a first simulation study we investigated the composition of a sample of all objects 3σ below the Ca II K index fit. For this we used a grid of model spectra with metallicities between $[\text{Fe}/\text{H}] = -1.0$ and -3.0 (Tab. 1). The model spectra have been converted to artificial objective-prism spectra by rebinning and multiplication with spectral sensitivity curves of the HES objective-prism plates. From Tab. 1 one can see that above $S/N = 15$ (corresponding to $B_J < 16.0 \dots 16.5$, depending on plate quality) there is a fairly sharp cutoff between $[\text{Fe}/\text{H}] = -1.0$ and $[\text{Fe}/\text{H}] = -2.0$, and the simulated samples of stars with $[\text{Fe}/\text{H}] \leq -2.0$ have a completeness close to 100 %. The restriction of the sample to lower metallicities is limited by the spectral resolution of the HES objective-prism spectra: For stars in the considered temperature range, Ca II K reaches the detection limit at $[\text{Fe}/\text{H}] \sim -2.0$.

A further test with *real* spectra of metal-poor stars confirms the high level of completeness, though the number of involved spectra is admittedly small. A cross-identification between 130 HES fields and the catalogue of BPS92 yielded 8 turnoff-stars with $[\text{Fe}/\text{H}] < -3.0$. All of these have been selected by the 3σ criterium.

3.2. Optimal Colour Range

The optimal colour range for an efficient selection of the most metal-poor, unevolved stars is $0.3 < (B - V) < 0.5$, since here no contamination of the sample with late-type giants is expected. The contamination with horizontal branch

Table 1. Selection function in dependence of stellar parameters T_{eff} , $\log g$, and $[\text{Fe}/\text{H}]$, at different magnitudes, for a selection by **CaKindex** $< 3\sigma$ below linear fit of **CaKindex** on one randomly choosen plate. For each class, 300 simulated model spectra have been used.

T_{eff}	$\log g$	$[\text{Fe}/\text{H}]$	$B - V$	% selected on randomly choosen plate					
				$S/N = 30$	25	20	15	10	5
				$B_J = 14.7$	15.2	15.7	16.1	16.7	17.3
6000	4.50	-1.00	0.50	0.0	0.0	0.0	0.0	20.0	37.3
6000	4.50	-2.00	0.48	98.3	96.7	91.3	81.0	98.7	84.3
6000	4.50	-3.00	0.47	100.0	100.0	99.7	99.7	100.0	93.7
6500	3.50	-1.00	0.39	0.0	0.0	0.0	0.0	43.3	43.0
6500	3.50	-2.00	0.37	96.3	92.0	88.7	83.0	99.0	82.7
6500	3.50	-3.00	0.37	100.0	99.3	97.0	92.3	98.7	91.3
6500	4.50	-1.00	0.41	0.0	0.0	0.0	0.3	30.7	41.0
6500	4.50	-2.00	0.40	97.3	96.0	90.0	79.3	98.7	86.3
6500	4.50	-3.00	0.40	100.0	100.0	97.7	96.0	100.0	90.0

stars will be small as well, since most of the colour range is overlapping with the RR Lyrae gap, i.e. $0.2 \lesssim (B - V) \lesssim 0.4$ (Walker 1998).

4. A First Set of New Metal-Poor Stars From the HES

In 1997 we selected a first set of candidates for metal-poor stars on HES plates by automatic classification, i.e., with a *different* method than described above. A learning sample of artificial, continous spectra was used. Follow-up observations were carried out with EMMI attached to the ESO-NTT at a spectral resolution of $\sim 5 \text{ \AA}$. Due to bad weather only six stars could be observed with a sufficient S/N ($\gtrsim 100$ at $\text{H}\alpha$) to determine their stellar parameters T_{eff} , $[\text{Fe}/\text{H}]$ and $\log g$ by spectral syntheses. The Balmer lines were used as a temperature indicator, Mg I b as a gravity indicator, and the Fe lines as a metallicity indicator. The results are given in Tab. 2.

Five of the six analysed stars are metal-poor, while one has $[\text{Fe}/\text{H}] = -0.4$. Given the experience we have gained at present, this star could have been easily rejected from the sample, because Mg I b is visible in the objective-prism spectrum.

5. Discussion

The above results suggest that for an efficient selection of the most metal-poor, unevolved stars we should restrict ourselves to objective-prism spectra with $S/N > 15$ in the colour range $0.3 < (B - V) < 0.5$. By applying the simulated selection function (Tab. 1) to the catalogue of BPS92, we estimate that at least $\sim 80\%$ of our candidates will be unevolved stars; $\sim 40\%$ of these having $[\text{Fe}/\text{H}] < -2.5$, and every tenth star having $[\text{Fe}/\text{H}] < -3.0$. A typical number of

Table 2. Stellar parameters of new metal poor stars discovered by the Hamburg/ESO survey. The spectrum of HE 0008–3842 shows inconsistent temperatures between different Balmer lines. For this star we list the parameters of the best possible fit.

Name	B_J	T_{eff}	$\log g$	[Fe/H]
HE 0131–2740	15.1	5600 ± 200 K	3.5 ± 0.5	-2.1 ± 0.5
HE 0008–3842	14.4	(4700 ± 300) K	(1.5 ± 0.5)	(-2.5 ± 0.5)
HE 0119–4211	15.6	5300 ± 200 K	2.0 ± 0.5	-2.3 ± 0.5
HE 0442–5113	15.5	5900 ± 200 K	3.7 ± 0.5	-1.9 ± 0.5
HE 0350–4804	16.4	6100 ± 200 K	4.5 ± 0.5	-2.0 ± 0.5
HE 0448–3524	15.3	6100 ± 200 K	3.8 ± 0.3	-0.4 ± 0.3

candidates is 50 per plate, and among them ~ 10 first-rank candidates. Therefore, even with the above restrictions applied, we will only be able to inspect our best candidates by single-slit spectroscopic follow-up observations. The most interesting of these will be investigated at high resolution using UVES and the VLT for age determinations and abundance studies.

An improved efficiency of our survey with respect to the yield of the most metal-poor stars is expected when automatic classification is used for the candidate selection, as more spectral features will be involved. An especially interesting feature is the G band of CH. Rossi et al. (1999, this volume) show that the most carbon-enhanced stars from the HK survey are found among the most metal-poor stars, so we expect that a selection of stars exhibiting strong G bands, and neither Mg I b nor Ca II K, will result in a sample of carbon-enhanced and extremely metal-poor stars.

Acknowledgments. N.C. was supported by DFG under grant Re 353/40-1 and travel grant Ch 214/1-1. T.C.B. acknowledges support for this work from AST 95-29454 awarded by the NSF.

References

- Beers, T.C., Preston, G.W., & Shectman, S.A. 1992, *AJ*, 103, 1987–2034
- Christlieb, N., Graßhoff, G., Nelke, A. & Wisotzki, L. 1997, in E. Kontizas et al. (eds.), *Wide-Field Spectroscopy*, 109–113
- Christlieb, N., Graßhoff, G., Nelke, A., Schlemminger, A. & Wisotzki, L. 1998a, in: I. Balderjahn et al. (eds.), *Classification, Data Analysis, and Data Highways*, 16–23
- Christlieb, N., Graßhoff, G., Nelke, A., Schlemminger, A. & Wisotzki, L. 1998b, in: R. Albrecht et al. (eds.), *Astronomical Data Analysis and Software Systems VII*, ASP Conference Series 145, 457–460
- Walker, A. R. 1998, *AJ*, 116, 220–236
- Watson, F. 1998, *AAO Newsletter*, 85, 11–12
- Wisotzki, L., Köhler, T., Groote, D. & Reimers, D. 1996, *A&AS*, 115, 227–233